

A scheme for dense coding in the non-symmetric quantum channel

Fengli Yan^{1,2} and Meiyu Wang²

¹ CCAST (World Laboratory), P.O. Box 8730, Beijing 100080, China

² Department of Physics, Hebei Normal University, Shijiazhuang 050016, China

(Dated: February 1, 2008)

We investigate the dense coding in the case of non-symmetric Hilbert spaces of the sender and receiver's particles sharing the quantum maximally entangled state. The efficiency of classical information gain is also considered. We conclude that when a more level particle is with the sender, she can get a non-symmetric quantum channel from a symmetric one by entanglement transfer. Thus the efficiency of information transmission is improved.

PACS numbers: 03.67.-a, 89.70.+c

I. INTRODUCTION

The quantum entanglement state [1, 2] is an important tool distinguishing the quantum mechanics from the classical physics. Due to the non-local properties of quantum entanglement state, it has been widely used in many fields such as quantum teleportation [3], quantum dense coding [4] and quantum key distribution [5]. Here we will focus our attention on quantum dense coding. In 1992, Bennett and Wiesner proposed the first scheme for the quantum dense coding [4], in which Bell states are used as quantum channel. In 1996, quantum dense coding was experimentally presented by Mattle *et al* in an optical system [6]. Recently Liu *et al* presented a protocol for dense coding with multi-level entangled states [7]. However in these schemes, the quantum channels are symmetric, that is to say, the dimension of the Hilbert space of the particle with sender is the same as that of the particle with receiver. In 1998, Bose *et al* generalized Bennett and Wiesner's scheme for dense coding into multi-parties, one receiver intends to receive messages from N senders [8]. In this case, the quantum channel is non-symmetric. The particles with senders are in 2^N -dimensional Hilbert space, while the particle with 2-dimensional Hilbert space belongs to receiver. In present paper, we give a scheme for quantum dense coding in the case of non-symmetric Hilbert spaces of the sender and receiver's particles sharing the quantum maximally entangled state, and consider the efficiency of classical information gain.

II. A SCHEME FOR DENSE CODING IN THE NON-SYMMETRIC QUANTUM CHANNEL

In Bennett and Wiesner's first scheme for the quantum dense coding, sender Alice and receiver Bob share a pair of entangled particles in the Bell state. Alice performs one of the four 1-qubit unitary operations given by the identity I or the Pauli matrices $(\sigma_x, i\sigma_y, \sigma_z)$ on her particle. Each of the unitary operations maps the Bell state to a different member of the four Bell states. Then Alice sends her particle to Bob. Bob can obtain two bits of classical information from the joint measurement on two

particles. Now we would like to generate the Bennett and Wiesner's symmetric scheme to the non-symmetric one.

To present our scheme clearly, let us first begin with dense coding between two parties in 3×2 -dimension. Suppose the particle 1 in 3-dimensional Hilbert space belongs to Alice, and the particle 2 in 2-dimensional Hilbert space is with Bob. They share the maximally entangled state:

$$|\Psi_{00}\rangle_{12} = \frac{1}{\sqrt{2}}(|00\rangle + |11\rangle)_{12}. \quad (1)$$

Through simple calculation, it can be shown that the single-body operators on particle 1:

$$\begin{aligned} U_{00} &= \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, & U_{01} &= \begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \\ U_{10} &= \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}, & U_{11} &= \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & -1 & 0 \end{bmatrix}, \\ U_{20} &= \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix}, & U_{21} &= \begin{bmatrix} 0 & -1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix} \end{aligned} \quad (2)$$

will transform $|\Psi_{00}\rangle$ into the corresponding state respectively:

$$U_{00}|\Psi_{00}\rangle = \frac{1}{\sqrt{2}}(|00\rangle + |11\rangle) = |\Psi_{00}\rangle, \quad (3)$$

$$U_{01}|\Psi_{00}\rangle = \frac{1}{\sqrt{2}}(|00\rangle - |11\rangle) = |\Psi_{01}\rangle, \quad (4)$$

$$U_{10}|\Psi_{00}\rangle = \frac{1}{\sqrt{2}}(|10\rangle + |21\rangle) = |\Psi_{10}\rangle, \quad (5)$$

$$U_{11}|\Psi_{00}\rangle = \frac{1}{\sqrt{2}}(|10\rangle - |21\rangle) = |\Psi_{11}\rangle, \quad (6)$$

$$U_{20}|\Psi_{00}\rangle = \frac{1}{\sqrt{2}}(|20\rangle + |01\rangle) = |\Psi_{20}\rangle, \quad (7)$$

$$U_{21}|\Psi_{00}\rangle = \frac{1}{\sqrt{2}}(|20\rangle - |01\rangle) = |\Psi_{21}\rangle. \quad (8)$$

Alice operates one of the above unitary transformations, and sends her particle to Bob. Bob takes only one measurement in the base $\{|\Psi_{00}\rangle, |\Psi_{01}\rangle, \dots, |\Psi_{21}\rangle\}$, and he will know what operation Alice has done, or to say, what the messages Alice has encoded in the quantum state. As a result, Bob gets $\log_2 6$ bits of information through only one measurement. Thus the dense coding is realized.

It is straightforward to generalize the above protocol to arbitrarily different dimension for two parties. Assume that the Hilbert space of Alice's particle 1 is non-symmetric with that of Bob's particle 2, i.e. the dimension p of the Hilbert space of particle 1 is not the same as the dimension q of particle 2. Without loss of generality, we choose $p > q$. The non-symmetric quantum channel shared by Alice and Bob is the maximally entangled state:

$$|\Psi_{00}\rangle_{12} = \frac{1}{\sqrt{q}}(|00\rangle + |11\rangle + \dots + |q-1, q-1\rangle)_{12}. \quad (9)$$

Clearly, an orthogonal base can be composed of the general Bell states of particle 1 and 2

$$|\Psi_{mn}\rangle = \sum_j e^{2\pi i j n / q} |(j \oplus m) \bmod p\rangle_1 \otimes |j\rangle_2 / \sqrt{q}, \quad (10)$$

where $m = 0, 1, \dots, p-1$; $n, j = 0, 1, \dots, q-1$. The single-body operators on particle 1 satisfying $U_{mn}|\Psi_{00}\rangle = |\Psi_{mn}\rangle$ may be explicitly written out:

$$U_{mn} = \sum_j e^{2\pi i j n / q} |(j \oplus m) \bmod p\rangle \langle j|, \quad (11)$$

where $m = 0, 1, \dots, p-1$; $n, j = 0, 1, \dots, q-1$. The procedure for realizing dense coding in this case is similar to the above one. In order to send information, Alice performs one of the operations in (11) to her particle and sends it to Bob. Bob then performs a collective measurement in the general Bell base $\{|\Psi_{mn}\rangle_{12} | m = 0, 1, \dots, p-1; n = 0, 1, \dots, q-1\}$ to find out what Alice has done to particle 1 and therefore reads out the encoded message. In this case, Bob can gain $\log_2 p \times q$ bits of information. If $p = q$, the quantum channel becomes symmetric, correspondingly the bits of information become $\log_2 p^2 = 2 \log_2 p$.

III. IMPROVING THE EFFICIENCY OF THE DENSE CODING BY ENTANGLEMENT TRANSFER

When the quantum channel between Alice and Bob is symmetric, besides a more level particle is with Alice, then the quantum channel may be transformed into non-symmetric one, and hence improve the efficiency of the

dense coding. For the sake of clearness, we start our discussion from the following simple case.

Suppose the initial quantum channel between Alice and Bob is

$$|\Phi\rangle_{12} = \frac{1}{\sqrt{2}}(|00\rangle + |11\rangle)_{12}, \quad (12)$$

the particle 1 belongs to Alice, and the particle 2 to Bob, they are both in 2-dimensional Hilbert space. Besides the particle 3 belonging to Alice is in 3-dimensional Hilbert space, its initial state is $|0\rangle_3$. So the initial state of the three particles is

$$|\Psi\rangle_{123} = \frac{1}{\sqrt{2}}(|00\rangle + |11\rangle)_{12} |0\rangle_3. \quad (13)$$

First Alice applies the following unitary operation on her particles 1 and 3 under the base $\{|00\rangle_{13}, |01\rangle_{13}, |02\rangle_{13}, |10\rangle_{13}, |11\rangle_{13}, |12\rangle_{13}\}$:

$$U_{13} = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}, \quad (14)$$

correspondingly $|\Psi\rangle_{123}$ becomes

$$U_{13}|\Psi\rangle_{123} = \frac{1}{\sqrt{2}}(|00\rangle_{13} |0\rangle_2 + |11\rangle_{13} |1\rangle_2). \quad (15)$$

Second Alice applies the following unitary operation on her particles 1 and 3:

$$U'_{13} = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}, \quad (16)$$

and hence equation (15) becomes

$$\begin{aligned} & U'_{13} \frac{1}{\sqrt{2}}(|00\rangle_{13} |0\rangle_2 + |11\rangle_{13} |1\rangle_2) \\ &= \frac{1}{\sqrt{2}}(|00\rangle_{23} + |11\rangle_{23}) |0\rangle_1 \\ &= |0\rangle_1 |\Phi'\rangle_{23}. \end{aligned} \quad (17)$$

As a result, the entanglement between the particles 1 and 2 is transferred to the entanglement between the particles 2 and 3. Before this, using the quantum channel between the particles 1 and 2 for dense coding, Bob gets $\log_2 4$ bits of information, but after that, Bob can gain $\log_2 6$ bits of information, so the efficiency of information transmission is increased.

In the following section, we consider the general case. Assume that Alice and Bob share the maximally entangled state:

$$|\Phi_{00}\rangle_{12} = \frac{1}{\sqrt{q}}(|00\rangle + |11\rangle + \dots + |q-1, q-1\rangle)_{12}, \quad (18)$$

where the particle 1 is with Alice, and the particle 2 with Bob, they are both in q -dimensional Hilbert space. Besides the particle 3 belonging to Alice is in p -dimensional Hilbert space, its initial state is $|0\rangle_3$, and $p > q$. Therefore three particles will be in the state $|\Phi_{00}\rangle_{12} \otimes |0\rangle_3$. Alice performs the following unitary operations on her particles 1 and 3:

$$U_{13} = \sum_{i=0}^{q-1} |ii\rangle\langle ii| + \sum_{j=1}^{q-1} |jj\rangle\langle jj| + \sum_{m,n} |mn\rangle\langle mn|, \quad (19)$$

where $m = 0, 1, \dots, q-1$; $n = 1, 2, \dots, p-1$, and $m \neq n$;

$$U'_{13} = \sum_{i=0}^{q-1} |0i\rangle\langle ii| + \sum_{j=1}^{q-1} |jj\rangle\langle 0j| + \sum_{m,n} |mn\rangle\langle mn|, \quad (20)$$

in the sum of above equation, when $m = 0$, we let $n = q, q+1, \dots, p-1$; when $m = 1, 2, \dots, q-1$, then n should take $n = 0, 1, \dots, p-1$ with the condition $m \neq n$. After that $|\Phi_{00}\rangle_{12} \otimes |0\rangle_3$ correspondingly becomes

$$\begin{aligned} & U'_{13} \otimes U_{13} |\Phi_{00}\rangle_{12} \otimes |0\rangle_3 \\ &= \frac{1}{\sqrt{q}} (|00\rangle + |11\rangle + \dots + |q-1, q-1\rangle)_{23} \otimes |0\rangle_1. \end{aligned} \quad (21)$$

So the entanglement transfer between the particles 1 and 2 and the particles 2 and 3 has been successfully realized.

In other words, a symmetric quantum channel has been transformed into the non-symmetric one. Clearly, before this, Bob gets $\log_2 q^2$ bits of information, now Bob can gain $\log_2 pq$ bits of information, and hence the efficiency of information transmission can be increased.

IV. CONCLUSION AND DISCUSSION

In summary, we have given a scheme to transform the symmetric level quantum maximally entangled state between two parties into non-symmetric one, and using it as the quantum channel for dense coding. We have also considered the efficiency of information transmission. Comparing our scheme with one for symmetric dense coding, obviously ours is more efficient. We hope this scheme can be realized by the experiment.

Acknowledgments

This work was supported by Hebei Natural Science Foundation under Grant No. 101094.

-
- [1] A. Einstein, B. Podolsky, and N. Rosen, Phys. Rev. **47**, 777 (1935).
 - [2] E. Schrödinger, Proc. Cambridge Philos. Soc. **31**, 555 (1935).
 - [3] C.H. Bennett, *et al*, Phys. Rev. Lett. **70**, 1895 (1993).
 - [4] C.H. Bennett and S.J. Wiesner, Phys. Rev. Lett. **69**, 2881 (1992).
 - [5] A.K. Ekert, Phys. Rev. Lett. **67**, 661 (1991).
 - [6] K. Mattle *et al*, Phys. Rev. Lett. **76**, 4656 (1996).
 - [7] X.S. Liu *et al*, Phys. Rev. **A** **65**, 022304 (2002).
 - [8] S. Bose, V. Vedral, and P.L. Knight, Phys. Rev. **A** **57**, 822 (1998).